

# On $\text{NS}_{\omega_1}$ being saturated

Ralf Schindler<sup>1</sup>

Institut für Mathematische Logik und Grundlagenforschung, Universität Münster  
Einsteinstr. 62, 48149 Münster, Germany

## 1 Preliminaries

Let  $\kappa$  be a regular uncountable cardinal. A normal uniform ideal  $I$  on  $\kappa$  is called *saturated* if forcing with  $I^+$ , ordered by  $S \leq T$  iff  $S \setminus T \in I$ , has the  $\kappa^+$ -c.c.  $I$  is called *presaturated* iff for all  $S \in I^+$  and for all maximal antichains  $A^n$ ,  $n < \omega$ , there is some  $T \leq S$ ,  $T \in I^+$ , such that for each  $n < \omega$ ,

$$\{R \in A^n : R \cap T \in I^+\}$$

has size at most  $\kappa$ . Finally,  $I$  is called *precipitous* iff the generic ultrapower is always well-founded. Every saturated ideal is presaturated, and every presaturated ideal is precipitous.

We shall produce two main theorems, due to S. Shelah (Theorem 2.1, see [4, Theorem 2.64]) and W.H. Woodin (Theorem 3.1, see [4, Theorem 2.61]) which say that a Woodin cardinal may be used to force  $\text{NS}_{\omega_1}$  to be saturated viz. presaturated. Theorem 2.3 is due to the current author. As it will be used in the proof of Theorem 3.1, we also give the construction of a model with the the Strong Chang Conjecture (Theorem 4.1).

**Definition 1.1** *Let  $\delta$  be a cardinal. We say that  $\delta$  is Woodin with  $\diamond$  iff there is some sequence  $(a_\kappa : \kappa < \delta)$  such that  $a_\kappa \subset V_\kappa$  for every  $\kappa < \delta$  and for every  $A \subset V_\delta$  the set*

$$\{\kappa < \delta : A \cap V_\kappa = a_\kappa \wedge \kappa \text{ is } A\text{-strong up to } \delta\}$$

*is stationary in  $\delta$ .*

**Lemma 1.2** *Suppose  $V = L[E]$ . Every Woodin cardinal is Woodin with  $\diamond$ .*

PROOF. Let us define  $((a_\kappa, c_\kappa) : \kappa < \delta)$  recursively as follows. If  $((a_\kappa, c_\kappa) : \kappa < \mu)$  is defined for some  $\mu < \delta$ , then we let  $(a_\mu, c_\mu)$  be the least (in the order of constructibility) pair  $(a, c)$  such that  $a \subset V_\mu$ ,  $c \subset \mu$  is club in  $\mu$ , and

$$\{\kappa < \mu : a \cap V_\kappa = a_\kappa \wedge \kappa \text{ is } a\text{-strong up to } \mu\} \cap c = \emptyset$$

(if such a pair  $(a, c)$  exists).

We claim that  $(a_\kappa : \kappa < \delta)$  is as desired. If not, then let  $(A, C)$  be least (in the order of constructibility) such that  $A \subset V_\delta$ ,  $C \subset \delta$  is club in  $\delta$ , and

$$(1) \quad \{\kappa < \delta : A \cap V_\kappa = a_\kappa \wedge \kappa \text{ is } A\text{-strong up to } \delta\} \cap C = \emptyset.$$

---

<sup>1</sup>The author thanks Daisuke Ikegami for helping him to figure out the details of the proof of Theorem 2.1 and for many other conversations on the topic of this note. He also thanks Andreas Lietz for helping him to figure out the details of the proof of Theorem 3.1. Written in Girona, Catalunya, Sept 05, 2011. An intermediate improved version as of April 24, 2020 was expanded significantly on Nov 13, 2021.

As the set

$$\{\kappa < \delta: \kappa \text{ is } A\text{-strong up to } \delta\}$$

is stationary in  $\delta$ , an easy Skolem hull argument together with condensation for  $L[E]$  yields some  $\kappa \in C$  which is  $A$ -strong up to  $\delta$  and  $(A \cap V_\kappa, c \cap \kappa)$  is the least (in the order of constructibility) pair  $(a, c)$  such that  $a \subset V_\kappa$ ,  $c \subset \kappa$  is club in  $\kappa$ , and

$$\{\lambda < \kappa: a \cap V_\lambda = a_\lambda \wedge \lambda \text{ is } a\text{-strong up to } \kappa\} \cap c = \emptyset.$$

But then  $(A \cap V_\kappa, c \cap \kappa) = (a_\kappa, c_\kappa)$ , which contradicts (1).  $\square$  (Lemma 1.2)

**Lemma 1.3** *Suppose that  $\delta$  is a Woodin cardinal. Then  $\delta$  is Woodin with  $\diamond$  in  $V^{\text{Col}(\delta, \delta)}$ .*

PROOF. We may identify  $\text{Col}(\delta, \delta)$  with the forcing

$$\mathbb{P} = \{(a_\kappa: \kappa < \mu): \mu < \delta \wedge \forall \kappa < \mu \ a_\kappa \subset V_\kappa\},$$

ordered by end-extension. Let  $\tau, \sigma \in V^\mathbb{P}$ , and let  $p \in \mathbb{P}$  be such that

$$p \Vdash \tau \subset V_\delta \wedge \sigma \subset \delta \text{ is club in } \delta.$$

We aim to find some  $q = (a_\lambda: \lambda < \mu) \leq p$  and some  $\kappa < \delta$  such that

$$q \Vdash \kappa \in \sigma \text{ is } \tau\text{-strong up to } \delta \wedge \tau \cap \kappa = a_\kappa.$$

Let us recursively construct a sequence  $(p_\kappa: \kappa < \delta) = ((a_\lambda: \lambda < \mu_\kappa))$  of stronger and stronger conditions end-extending  $p$  with the following properties.

- (a)  $\{\mu_\kappa: \kappa < \delta\}$  is club in  $\delta$ .
- (b) For all  $\kappa$  there is some  $c_\kappa \subset \mu_\kappa$  which is unbounded in  $\mu_\kappa$  such that  $p_\kappa \Vdash \sigma \cap \mu_\kappa = c_\kappa$ ; in particular,  $p_\kappa \Vdash \mu_\kappa \in \sigma$ .
- (c) For all  $\kappa$  there is some  $A_\kappa \subset V_{\mu_\kappa}$  such that  $p_\kappa \Vdash \tau \cap V_{\mu_\kappa} = A_\kappa$ .
- (d) For all  $\kappa$ ,  $a_{\mu_\kappa} = A_\kappa$ .
- (e) If  $(a_\lambda: \lambda < \mu_{\kappa+1})$  does not force  $\kappa$  to be  $\tau$ -strong up to  $\delta$ , then there is some  $\alpha < \mu_{\kappa+1}$  such that

$$p_{\kappa+1} \Vdash \kappa \text{ is not } \tau\text{-strong up to } \alpha.$$

There is no problem with this construction.

Now set  $A = \bigcup_{\kappa < \delta} A_\kappa$ , so that  $A \cap V_{\mu_\kappa} = A_\kappa$  for all  $\kappa$ . As  $\delta$  is Woodin, by (a) we may pick some  $\kappa = \mu_\kappa$  which is  $A$ -strong up to  $\delta$ . Set  $q = (a_\lambda: \lambda < \kappa + 1)$ . By (b), (c), (d) we have that

$$q \Vdash \kappa \in \sigma \wedge \tau \cap \kappa = a_\kappa.$$

If  $q$  does not force  $\kappa$  to be  $\tau$ -strong up to  $\delta$ , then by (c), (e), and the definition of  $A$ , there is some  $\alpha < \mu_{\kappa+1}$  with

$$p_{\kappa+1} \Vdash \kappa \text{ is not } A\text{-strong up to } \alpha,$$

which is nonsense.

$q$  is thus as desired.  $\square$  (Lemma 1.3)

## 2 Forcing $\text{NS}_{\omega_1}$ to be saturated

**Theorem 2.1 (Shelah)** *Let  $\delta$  be a Woodin cardinal. There is some semi-proper  $\mathbb{P} \subset V_\delta$  with the  $\delta$ -c.c. such that if  $G$  is  $\mathbb{P}$ -generic over  $V$ , then  $V[G] \models$  “ $\text{NS}_{\omega_1}$  is saturated.”*

PROOF. Let us assume that  $\delta$  is Woodin with  $\diamond$ . We perform an RCS iteration (cf. [1]) of length  $\delta + 1$  of semi-proper forcings each of size  $< \delta$ , where in each successor step of the iteration, we either force with the poset  $\mathbb{S}(\vec{S})$  to seal a given maximal antichain  $\vec{S} \subset (\text{NS}_{\omega_1})^+/\text{NS}_{\omega_1}$ , provided that  $\mathbb{S}(\vec{S})$  is semi-proper, or else we force with  $\text{Col}(\omega_1, 2^{\aleph_2})$  (which is  $\omega$ -closed, hence [semi-]proper). The choice of the maximal antichain  $\vec{S}$  is according to the  $\diamond$ -Woodinness of  $\delta$  and will be left to the reader's discretion.

If  $\vec{S}$  is a (not necessarily maximal) antichain, then the sealing forcing  $\mathbb{S}(\vec{S})$  consists of all pairs  $(c, p)$  such that for some  $\beta < \omega_1$  we have that  $c: \beta + 1 \rightarrow \omega_1$ ,  $p: \beta + 1 \rightarrow \vec{S}$ ,  $\text{ran}(c)$  is a closed subset of  $\omega_1$ , and for all  $\xi \leq \beta$ ,  $c(\xi) \in \bigcup_{i < \xi} p(i)$ .  $\mathbb{S}(\vec{S})$  is ordered by end-extension. The forcing  $\mathbb{S}(\vec{S})$  is  $\omega$ -distributive and preserves all the stationary subsets of all  $S \in \vec{S}$ , so that  $\mathbb{S}(\vec{S})$  is stationary set preserving if  $\vec{S}$  is maximal.

Let us write  $\mathbb{P}$  for the entire iteration. Let us pick some  $G$  which is  $\mathbb{P}$ -generic over  $V$ . We aim to prove that in  $V[G]$ , every antichain in  $(\text{NS}_{\omega_1})^+/\text{NS}_{\omega_1}$  has size  $\leq \aleph_1$ .

Suppose not, and let  $\vec{S} = (S_i: i < \delta) \in V[G]$  be a maximal antichain. Let  $\vec{S} = \tau^G$ , where  $\tau \in V^{\mathbb{P}} \cap V_{\delta+1}$ . We may find some  $\kappa < \delta$  such that

- (i)  $\kappa$  is  $\mathbb{P} \oplus \tau$ -strong up to  $\delta$  in  $V$ ,
- (ii)  $\kappa = \omega_2^{V[G \upharpoonright \kappa]}$ , and
- (iii)  $\vec{S} \upharpoonright \kappa = (S_i: i < \kappa) = (\tau \cap V_\kappa)^{G \upharpoonright \kappa}$  is the maximal antichain in  $V[G \upharpoonright \kappa]$  which is picked at stage  $\kappa$ .

The forcing  $\mathbb{S}(\vec{S} \upharpoonright \kappa)$  for sealing  $\vec{S} \upharpoonright \kappa$ , as defined in  $V[G \upharpoonright \kappa]$ , cannot be semi-proper in  $V[G \upharpoonright \kappa]$ , so that there is some  $(c, p) \in \mathbb{S}(\vec{S} \upharpoonright \kappa)$  such that the set

$$\begin{aligned} \tilde{T} = \{ & X \prec (H_{\kappa^+})^{V[G \upharpoonright \kappa]} : \text{Card}(X) = \aleph_0 \wedge (c, p) \in X \wedge \neg \exists Y \supset X (Y \prec (H_{\kappa^+})^{V[G \upharpoonright \kappa]} \wedge \\ & \text{Card}(Y) = \aleph_0 \wedge Y \cap \omega_1 = X \cap \omega_1 \wedge \exists (d, q) \leq (c, p) \quad (d, q) \text{ is } Y\text{-generic} ) \} \end{aligned}$$

is stationary in  $V[G \upharpoonright \kappa]$ , and the  $\kappa^{\text{th}}$  forcing in the iteration  $\mathbb{P}$  is  $\text{Col}(\omega_1, 2^{\aleph_2})$ . In  $V[G \upharpoonright \kappa + 1]$  there is a surjective  $f: \omega_1 \rightarrow (H_{\kappa^+})^{V[G \upharpoonright \kappa]}$ . Because  $\text{Col}(\omega_1, 2^{\aleph_2})$  is proper,  $\tilde{T}$  is still stationary in  $V[G \upharpoonright \kappa + 1]$ , and hence the set

$$T = \{ \alpha < \omega_1 : f'' \alpha \in \tilde{T} \wedge \alpha = f''' \alpha \cap \omega_1 \}$$

is stationary in  $V[G \upharpoonright \kappa + 1]$ . As the tail  $\mathbb{P}_{[\kappa+1, \delta]}$  of the iteration  $\mathbb{P}$  over  $V[G \upharpoonright \kappa + 1]$  is semi-proper,  $T$  will remain stationary in  $V[G]$ , and as  $\vec{S}$  is a maximal antichain there is some (unique)  $i_0 < \delta$  such that

- (2)  $T \cap S_{i_0}$  is stationary in  $V[G]$ .

(It is not hard to verify that  $i_0 \geq \kappa$ .)

Let  $\lambda < \delta$ ,  $\lambda > \max(i_0, \kappa + 1)$  be such that  $(\tau \cap V_\lambda)^{G \upharpoonright \lambda} = \bar{S} \upharpoonright \lambda$ , so that  $S_{i_0} = (\tau \cap V_\lambda)^{G \upharpoonright \lambda}(i_0)$ , the  $(i_0)^{\text{th}}$  element of  $(\tau \cap V_\lambda)^{G \upharpoonright \lambda}$ . Pick an elementary embedding

$$j: V \rightarrow M$$

such that  $\text{crit}(j) = \kappa$ ,  $M$  is transitive,  ${}^\kappa M \subset M$ ,  $V_{\lambda+\omega} \subset M$ ,  $j(\mathbb{P}) \cap V_\lambda = \mathbb{P} \cap V_\lambda$ , and  $j(\tau) \cap V_\lambda = \tau \cap V_\lambda$ .

Let  $(X_i: i < \omega_1) \in V[G \upharpoonright \kappa + 1]$  be an increasing continuous chain of countable substructures of  $(H_{j((2^\kappa)^+)})^{M[G \upharpoonright \kappa + 1]}$  with  $\{\tau \cap V_\lambda, i_0\} \subset X_0$  and such that for all  $i < \omega_1$ ,

- (a)  $i \in X_{i+1}$ ,
- (b)  $f''(X_i \cap \omega_1) \subset X_i$ , and
- (c)  $j''(X_i \cap (2^\kappa)^{V[G \upharpoonright \kappa]}) \subset X_i$ .

Write  $\bar{G} = G \upharpoonright [\kappa + 2, \lambda]$ . We have that

$$\{X_i[\bar{G}] \cap \omega_1: i < \omega_1\} \in V[G \upharpoonright \lambda]$$

is club in  $\omega_1$ , so that by (2) we may find some  $i < \omega_1$  with  $X_i[\bar{G}] \cap \omega_1 = X_i \cap \omega_1 \in T \cap S_{i_0}$ .

Write  $X = X_i$  and  $\alpha = X \cap \omega_1$ . As  $\text{Col}(\omega_1, 2^{\aleph_2})$  is  $\omega$ -closed,  $X \cap (H_{\kappa^+})^{V[G \upharpoonright \kappa]} \in V[G \upharpoonright \kappa]$ . As  $\alpha \in T$ ,  $f''\alpha \in \tilde{T}$  and  $\alpha = f''\alpha \cap \omega_1$ , and hence by (b)

$$f''\alpha \subset X \cap (H_{\kappa^+})^{V[G \upharpoonright \kappa]} \in V[G \upharpoonright \kappa].$$

This implies that

$$(3) \quad X \cap (H_{\kappa^+})^{V[G \upharpoonright \kappa]} \in \tilde{T}.$$

As the segment  $(\mathbb{P}_{[\lambda+1, j(\kappa)]})^{M[G \upharpoonright \lambda]}$  of  $j(\mathbb{P})$  over  $M[G \upharpoonright \lambda]$  is semi-proper, we may now pick  $H$  to be generic for the segment  $(\mathbb{P}_{[\lambda+1, j(\kappa)]})^{M[G \upharpoonright \lambda]}$  of  $j(\mathbb{P})$  over  $M[G \upharpoonright \lambda]$  in such a way that

$$X[\bar{G}, H] \cap \omega_1 = X[\bar{G}] \cap \omega_1.$$

We may lift  $j: V \rightarrow M$  to an elementary embedding

$$j^*: V[G \upharpoonright \kappa] \rightarrow M[G \upharpoonright \lambda, H].$$

Notice that  $(V_{\lambda+\omega})^{M[G \upharpoonright \lambda]} = (V_{\lambda+\omega})^{V[G \upharpoonright \lambda]}$ . By (3),

$$(4) \quad j^*(X \cap (H_{\kappa^+})^{V[G \upharpoonright \kappa]}) \in j^*(\tilde{T}).$$

We now have that  $X[\bar{G}, H] \cap \omega_1 = X[\bar{G}] \cap \omega_1 = \alpha \in S_{i_0} = (\tau \cap V_\lambda)^{G \upharpoonright \lambda}(i_0) \in X[\bar{G}] \subset X[\bar{G}, H] \prec (H_{j((2^\kappa)^+)})^{M[G \upharpoonright \lambda, H]}$ .

But now by (c),

$$j^*(X \cap (H_{\kappa^+})^{V[G \upharpoonright \kappa]}) = j^{**}(X \cap (H_{\kappa^+})^{V[G \upharpoonright \kappa]}) \subset X[\bar{G}, H].$$

Therefore,  $X[\bar{G}, H]$  witnesses that  $j^*(X \cap (H_{\kappa^+})^{V[G \upharpoonright \kappa]})$  is *not* in  $j^*(\bar{T})$ , as the condition  $j((c, p)) = (c, p) \in \mathbb{S}(\bar{S} \upharpoonright \kappa) \subset j(\mathbb{S}(\bar{S} \upharpoonright \kappa))$  from the definition of  $\bar{T}$  may be extended in  $j(\mathbb{S}(\bar{S} \upharpoonright \kappa))$  to some  $X[\bar{G}, H]$ -generic condition  $(c^*, p^*) \in j(\mathbb{S}(\bar{S} \upharpoonright \kappa))$  with  $\text{dom}(c^*) = \text{dom}(p^*) = \alpha + 1$ ,  $c^*(\alpha) = \alpha$ , and  $p^*(i) = S_{i_0} \in j^*(S \upharpoonright \kappa)$  for some  $i < \alpha$ .

This contradicts (4). □ (Theorem 2.1)

**Theorem 2.2 (Woodin)** *Suppose that  $\text{NS}_{\omega_1}$  is saturated and  $(\mathcal{P}(\omega_1))^\#$  exists. Then  $\delta_2^1 = \omega_2$ .*

PROOF SKETCH. (Cf. [4].) If  $N \cong X \prec \mathcal{M} = ((\mathcal{P}(\omega_1))^\#; \in, \text{NS}_{\omega_1})$ , where  $N$  is countable and transitive, then  $N$  is generically  $(\omega_1 + 1)$ -iterable via the preimage of  $\text{NS}_{\omega_1}$  and its images. By the Boundedness Lemma, the ordinal height of every  $(\omega_1)^{\text{th}}$  iterate of  $N$  is  $< (\omega_1^V)^{+L[z]}$ , where  $z \in \mathbb{R}$  codes  $N$ . On the other hand, if  $N_i \cong X_i = \text{Hull}^{\mathcal{M}}(X \cup \{X_j \cap \omega_1 : j < i\}) \prec \mathcal{M}$  for  $i \leq \omega_1$ , then  $(N_i : i \leq \omega_1)$ , together with the obvious maps, is a generic iteration of  $N$ . Hence if  $\beta \in X$ , where  $\beta < \omega_2$ ,  $\beta < (\omega_1^V)^{+L[z]} < \delta_2^1$ . □ (Theorem 2.2)

[4] shows that if  $\mathbb{P}$  is the poset of Theorem 2.1, as defined over  $M_1$ , and if  $G$  is  $\mathbb{P}$ -generic over  $M_1$ , then  $\delta_2^1 < \omega_2$  in  $M_1[G]$ . The following Theorem gives a bit more information.

**Theorem 2.3** *Let  $\mathbb{P}$  be the poset of Theorem 2.1, as defined over  $M_1$ , and let  $G$  be  $\mathbb{P}$ -generic over  $M_1$ . Then  $(\delta_2^1)^{M_1[G]} = (\delta_2^1)^{M_1} < \omega_2^{M_1} < \omega_2^{M_1[G]}$ .*

PROOF. Deny. Let  $x \in \mathbb{R} \cap M_1[G]$  witness that  $(\delta_2^1)^{M_1[G]} > (\delta_2^1)^{M_1}$ . So if

$$(N_i, \pi_{ij} : i \leq j \leq \omega_1)$$

is the iteration of  $x^\dagger = N_0$  of length  $\omega_1 + 1$  which is obtained by hitting the bottom (total) measure of  $x^\dagger$  and its images  $\omega_1$  times, then  $(\omega_1^V)^{+N_{\omega_1}} > (\delta_2^1)^{M_1}$ .

As  $x^\dagger \models$  “There is no inner model with a Woodin cardinal,” we may let  $K$  denote the core model of  $x^\dagger$  of height  $\Omega$ , where  $\Omega$  is the top measurable cardinal of  $x^\dagger$ . By [3], there is a normal iteration tree  $\mathcal{T}$  on  $K$  with  $[0, \infty)_{\mathcal{T}} = \emptyset$  and last model  $K^{N_1}$  such that  $\pi_{01} = \pi_{0\infty}^{\mathcal{T}}$ . Letting  $\mathcal{T}^*$  be the concatenation of all  $\pi_{0i}(\mathcal{T})$ ,  $0 \leq i < \omega_1$ ,  $\mathcal{T}^*$  is then a (non-normal) iteration tree on  $K$  with  $[0, \infty)_{\mathcal{T}^*} = \emptyset$  and last model  $K^{N_{\omega_1}}$  such that  $\pi_{0\omega_1} \upharpoonright K = \pi_{0\infty}^{\mathcal{T}^*}$ . By absoluteness,  $K$  is in fact iterable in  $M_1[G]$ , and  $\mathcal{T}^*$  is according to the (unique) relevant iteration strategy.

We claim that  $K$  iterates past  $M_1 \upharpoonright \omega_1$ .

Otherwise suppose that  $\alpha < \omega_1$  is such that  $M_1 \upharpoonright \alpha$  absorbs  $K$ . There is then, in  $M_1[G]$ , an iteration tree  $\mathcal{U}$  on  $M_1 \upharpoonright \alpha$  of length  $\omega_1 + 1$  such that  $\mathcal{M}_{\omega_1}^{\mathcal{U}} \cap \text{OR} \geq N_{\omega_1} \cap \text{OR} > (\delta_2^1)^{M_1}$ . (Cf. [2] for a writeup of this argument.) On the other hand, by the Boundedness Lemma, if  $z \in \mathbb{R} \cap M_1$  codes  $M_1 \upharpoonright \alpha$  and if  $\gamma$  denotes the supremum of all the ordinal heights of all  $(\omega_1)^{\text{th}}$  iterates of  $M_1 \upharpoonright \alpha$ , then

$$\gamma < (\omega_1)^{+L[z]}.$$

In particular,  $(\delta_2^1)^{M_1} > (\omega_1)^{+L[z]} > \gamma > \mathcal{M}_{\omega_1}^{\mathcal{U}} \cap \text{OR} > (\delta_2^1)^{M_1}$ .

This contradiction indeed shows that  $K$  iterates past  $M_1 \upharpoonright \omega_1$ . But then  $\omega_1$  has to be an inaccessible cardinal of  $M_1$ , which is nonsense. □ (Theorem 2.3)

**Question 1.** Is it true that  $M_1[G] \models \neg\text{CH}$  ?

**Question 2.** Is it true that  $\mathbb{R} \cap M_1[G] \subset M_1$  ?

### 3 Forcing $\text{NS}_{\omega_1}$ to be presaturated

**Theorem 3.1** *Let  $\delta$  be a Woodin cardinal. If  $G$  is  $\text{Col}(\omega_1, < \delta)$ -generic over  $V$ , then  $V[G] \models$  “ $\text{NS}_{\omega_1}$  is presaturated.”*

PROOF. Inside  $V[G]$ , let  $S \subset \omega_1$  be stationary, and let  $(A^n : n < \omega)$  be a sequence of maximal antichains in  $\text{NS}_{\omega_1}^+$ . By the argument from the proof of Theorem 2.1 we may find some  $\kappa < \delta$  which in  $V$  is sufficiently strong and such  $\{S, (A^n \cap V[G \upharpoonright \kappa] : n < \omega)\} \subset V[G \upharpoonright \kappa]$  and that in  $V[G \upharpoonright \kappa]$  for every  $n < \omega$  the set

$$\tilde{T}^n = \{X \prec (H_{\kappa^+})^{V[G \upharpoonright \kappa]} : \text{Card}(X) = \aleph_0 \wedge \exists Y \supset X (Y \prec (H_{\kappa^+})^{V[G \upharpoonright \kappa]} \wedge$$

$$\text{Card}(Y) = \aleph_0 \wedge Y \cap \omega_1 = X \cap \omega_1 \wedge \exists S \in A^n \cap V[G \upharpoonright \kappa] (\omega_1 \cap X \in S \in Y)\}$$

is stationary.  $V[G \upharpoonright \kappa]$  also satisfies the Strong Chang Conjecture, see the Appendix and specifically Theorem 4.1. We may then work in  $V[G \upharpoonright \kappa]$  and find some  $Y \prec (H_{\kappa^+})^{V[G \upharpoonright \kappa]}$  such that

- (a)  $\text{Card}(Y) = \aleph_0$ ,  $Y \cap \omega_1 \in S$ ,
- (b) for all  $n < \omega$  there is some  $S \in A^n \cap V[G \upharpoonright \kappa]$  with  $\omega_1 \cap Y \in S \in Y$ , and
- (c) there is some  $A \in Y$  of size  $\aleph_1$  such that for each  $n < \omega$  there is some  $S \in A$  as in (b).

Write  $\alpha = Y \cap \omega_1$ . Let  $f : \omega_1 \rightarrow A$  be onto,  $f \in Y$ . For  $n < \omega$ , let

$$T^n = \{\xi < \omega_1 : \exists \eta < \xi \ \xi \in f(\eta) \in A^n \cap V[G \upharpoonright \kappa]\}.$$

By (c),  $\alpha \in T^n$  for each  $n < \omega$ . Therefore,

$$(5) \quad \alpha \in S \cap \bigcap_{n < \omega} T^n.$$

As  $\{S, (T^n : n < \omega)\} \subset Y$ , (5) shows that  $T = S \cap \bigcap_{n < \omega} T^n$  is stationary in  $V[G \upharpoonright \kappa]$  and therefore in  $V[G]$ .

Let  $n < \omega$ . We claim that if  $R \in A^n \setminus A$ , then  $R \cap T$  is nonstationary. For suppose that  $R \cap T$  were stationary, so that  $R \cap T^n$  would also be stationary. For  $\xi \in R \cap T^n$ , pick  $\eta = \eta(\xi) < \xi$  such that  $\xi \in f(\eta) \in A^n \cap V[G \upharpoonright \kappa]$ . There is some stationary  $W \subset R \cap T^n$  and some  $\eta_0$  such that  $\eta_0 = \eta(\xi)$  for all  $\xi \in W$ , i.e.,  $W \subset f(\eta_0) \in A^n \cap A$ . But then  $R \cap f(\eta_0)$  would be stationary and  $A^n$  would not be an antichain.  $\square$  (Theorem 3.1)

## 4 Appendix: The Strong Chang Conjecture

We say that the *Strong Chang Conjecture* holds true if for every sufficiently big  $\theta$ , if  $X \prec H_\theta$  is countable, then there is some  $Y \prec H_\theta$  such that  $X \subset Y$ ,  $Y \cap \omega_1 = X \cap \omega_1$ , and  $\sup(X \cap \omega_2) < \sup(Y \cap \omega_2)$ .

**Theorem 4.1** *Let  $\kappa$  be a measurable cardinal, and let  $g$  be  $\text{Col}(\omega_1, < \kappa)$ -generic over  $V$ . The Strong Chang Conjecture holds true in  $V[g]$ .*

PROOF. Let  $\theta$  be sufficiently big, and let  $X \prec (H_\theta)^{V[g]}$  be countable. Write  $\alpha = X \cap \omega_1$ . Let  $\sigma: N \cong X$ , where  $N$  is transitive. We may write  $N = \bar{N}[\bar{g}]$ , and we have  $\sigma \upharpoonright \bar{N}: \bar{N} \rightarrow (H_\theta)^V$  is elementary,  $\bar{g} = \sigma^{-1}(g)$ , and  $\{\bar{N}, \bar{g}, \sigma \upharpoonright \bar{N}, \sigma''\bar{g}\} \subset V$ .

Let  $\bar{\kappa} = \sigma^{-1}(\kappa)$ , let  $U \in \text{ran}(\sigma)$  be a measure on  $\kappa$ , and let  $\bar{U} = \sigma^{-1}(U)$ . Let

$$i: \bar{N} \rightarrow P = \text{ult}(\bar{N}; \bar{U})$$

be the ultrapower map, where  $P$  is transitive.

By standard arguments, there is some  $Z \in U$  such that for every  $\beta \in Z$ , the map

$$\sigma^\beta: i(f)(\bar{\kappa}) \mapsto \sigma(f)(\beta)$$

defines an elementary embedding  $\sigma^\beta: P \rightarrow (H_\theta)^V$  with  $\sigma^\beta \circ i = \sigma$ .

Let  $h$  be  $\text{Col}(\alpha, [\bar{\kappa}, i(\bar{\kappa})])^P$ -generic over  $P[\bar{g}]$ ,  $h \in V$ . For  $\beta \in Z$ ,  $\sigma^{\beta''}h \in V$ , and also

$$q_\beta = \bigcup \sigma^{\beta''}h \in \text{Col}(\omega_1, [\beta, \kappa]),$$

so that we may construe  $q_\beta$  as an element of  $\text{Col}(\omega_1, < \kappa)$ .

As the support of  $q_\beta$  is a (countable) subset of  $[\beta, \kappa)$  and  $Z$  is unbounded in  $\kappa$ , an easy argument shows that

$$\{p \in \text{Col}(\omega_1, < \kappa) : \exists \beta \in Z p \leq q_\beta\}$$

is dense in  $\text{Col}(\omega_1, < \kappa)$ . Therefore there is some  $\beta \in Z$  such that  $q_\beta \in g$ .

For such  $\beta$ , we may extend  $\sigma^\beta$  to an map  $\hat{\sigma}^\beta: P[\bar{g}, h] \rightarrow (H_\theta)^{V[g]}$  defined by

$$\hat{\sigma}^\beta: \tau^{\bar{g}, h} \mapsto (\sigma^\beta(\tau))^g.$$

As  $\sigma^{\beta''}\bar{g}, h \subset g$ ,  $\hat{\sigma}^\beta$  is well-defined and elementary.

But then  $Y = \text{ran}(\hat{\sigma}^\beta)$  is as desired.

□ (Theorem 4.1)

## References

- [1] Ulrich Fuchs, *Donder's Version of Revised Countable Support*, available at <https://arxiv.org/abs/math/9207204>
- [2] Ikegami, D., *Ph.D. Thesis*, University of Amsterdam, available at <http://www.sic.shibaura-it.ac.jp/~ikegami/thesis-latest.pdf>
- [3] Schindler, R., *Iterates of the core model*, 71 (2006), pp.241-251.
- [4] Woodin, H.W., *The axiom of determinacy, forcing axioms, and the non-stationary ideal*, Berlin, New York 1999.